

## Anti-PRKAB1 antibody (100-200) (STJ110230)

ST.1110230

## **GENERAL INFORMATION**

Product Type Primary antibodies

**Short Description** 

Applications WB/IHC-P/ELISA Host/Source Rabbit

Reactivity Human/Mouse/Rat

## **PRODUCT PROPERTIES**

Clonality Polyclonal

Clone ID
Concentration
Conjugation
Purification
Dilution Range
WB:1:500-1:2000

IHC-P:1:50-1:200

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.

**Isotype** IgG

Storage Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

Instruction

## **TARGET INFORMATION**

Gene ID 5564
Gene Symbol PRKAB1
Uniprot ID AAKB1\_HUMAN


Immunogen Immunogen 100-200

mmunogen 100-20 Region


**Specificity** A synthetic peptide corresponding to a sequence within amino acids 100-200 of human AMPK Beta 1 (NP\_006244.2).

Immunogen WSKLPLTRSHNNFVAILDLP EGEHQYKFFVDGQWTHDPSE PIVTSQLGTVNNIIQVKKTD FEVFDALMVDSQKCSDVSEL


Sequence SSSPPGPYHQEPYVCKPEER F




Western blot analysis of extracts of various cell line using AMPK Beta 1 antibody (STJ110230) at 1:100 dilution. Secondary antibody: HRP Goat Anti-rabbit Ig (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per line 11 to 1000 dilution. Lysates/proteins: 25ug per line 11 to 1000 dilution.



Immunohistochemistry analysis of paraffin-embedde mouse kidney using AMPK Beta 1 artiblo (STJ110230) at dilution of 1:100 (40x lens). Perfor microwave antigen retrieval with 10 mM PBS buffer p 7. 2 before commencing with immunohistochemist statistics protocol



Immunohistochemistry of paraffin-embedded rat spleen using AMPK Beta 1 antibody (STJ110230) at dilution of 1:100 (40x lens).



Immunohistochemistry analysis of paraffin-embedded rat kidney using AMPK Beta 1 antibody (STJ110230) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commercing with immunohistochemistry staining