

Anti-NAPA antibody (1-295) (STJ110255)

STJ110255

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB:1:500-1:2000
Range	IHC-P:1:100-1:200
	ELISA:Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	8775
Gene Symbol	NAPA
Uniprot ID	SNAA_HUMAN
Immunogen	
Immunogen Region	1-295
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-295 of human Alpha SNAP (NP_003818.2).
Immunogen Sequence	MDNSGKEAEAMALLAEAEERK VKNSQSFFSGLFGSSKIEE ACEIYARAANMFMAKNWSA AGNAFCQAAQLHLQLQSKHD AATCFVDAGNAFKKADPQEA INCLMRAIEIYDMGRFTIA AKHHISIAEIYETELVDIEK AIAHYEQSADYKGEESNSS ANKCLLKVAGYAALLEQYQK AIDIYEQVGTNAMDSPLLY SAKDYFFKAALCHFCIDMLN AKLAVQKYEELFPAFSDSR




Western blot analysis of extracts of various cell lines, using Alpha SNAP antibody (STJ110255) at 1:500 dilution. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 90s.



Immunohistochemistry analysis of paraffin-embedded human prostate using Alpha SNAP antibody (STJ110255) (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry of paraffin-embedded mouse brain using Alpha SNAP antibody (STJ110255) (40x lens).



Immunohistochemistry analysis of paraffin-embedded rat brain using Alpha SNAP antibody (STJ110255) (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081