

## Anti-BLOC1S2 antibody (44-142) (STJ110314)

STJ110314

## **GENERAL INFORMATION**

Product Type Primary antibodies

**Short Description** 

Applications WB/ELISA Host/Source Rabbit

Reactivity Human/Mouse/Rat

## **PRODUCT PROPERTIES**

Clonality Polyclonal

Clone ID
Concentration Lot specific
Conjugation Unconjugated
Purification Affinity purification
Dilution Range WB:1:500-1:2000

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.

Isotype IgG

Storage Instruction Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

## **TARGET INFORMATION**

Gene ID 282991
Gene Symbol BLOC1S2
Uniprot ID BL1S2 HU


Uniprot ID BL1S2\_HUMAN Immunogen

Immunogen Region 44-142


**Specificity** Recombinant fusion protein containing a sequence corresponding to amino acids 44-142 of human BLOC1S2 (NP\_776170.2).

Immunogen MFSKMATYLTGELTATSEDY KLLENMNKLTSLKYLEMKDI AINISRNLKDLNQKYAGLQP YLDQINVIEEQVAALEQAAY

Sequence KLDAYSKKLEAKYKKLEKR



Nestern blot analysis of extracts of various cell line: using BLOC152 antibody (STJ1110314) at 1:100 dilution. Secondary antibody: HRP Goat Anti-rabling (H-LI) at 1:10000 dilution. Lysates/proteins: 25u per lane. Blocking buffer: 3% non-fat dry milk in TBS Detection: ECL Enhanced Kit. Exposure time: 30s.



Western blot analysis of extracts of various cell lines using BLOC152 antibody (STJ110314) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbi IgG (H+L) (STJS000856) at 1:10000 dilution Lysates/profisirs: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit Evropsure time: 30s