

**Anti-DHX58 antibody (389-678) (STJ110556)**  
STJ110556

**GENERAL INFORMATION**


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	
<b>Applications</b>	WB/IF/ICC/ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human/Mouse/Rat

**PRODUCT PROPERTIES**


<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	Lot specific
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purification
<b>Dilution Range</b>	WB: 1:500-1:2000 IF/CC: 1:50-1:200 ELISA: Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
<b>Formulation</b>	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
<b>Isotype</b>	IgG
<b>Storage Instruction</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

**TARGET INFORMATION**


<b>Gene ID</b>	79132
<b>Gene Symbol</b>	DHX58
<b>Uniprot ID</b>	DHX58_HUMAN
<b>Immunogen</b>	
<b>Immunogen Region</b>	389-678
<b>Specificity</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 389-678 of human DHX58 (NP_077024.2).
<b>Immunogen Sequence</b>	QQQGLQTVDIRAQLLIGAGN SSQSTHMTQRDQQEVIQKFQ DGTLLNLLVATSVAEGLDIP HCNVVVRYGLLTNEISMVQA RGRARADQSVYAFVATEGSR ELKRELINEAETLMEQAVA AVQKMDQAEYQAKIRDLQQA ALTKRAAQAAQRENQRQQFP VEHVQLLINCINMVAVGHGSD LRVVEGTHHVNPNFNSNY NVSRDPVWINKVFKDWKPGG VISCRCGEVWGLQMIYKS




Western blot analysis of extracts of various cell lines, using DHX58 antibody (STJ110556) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 60s.



Immunofluorescence analysis of U2OS cells using DHX58 antibody (STJ110556) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using DHX58 antibody (STJ110556) at dilution of 1:100. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates, using DHX58 Rabbit polyclonal antibody (STJ110556) at 1:9000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ5000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 8081