

Anti-RIPK4 antibody (240-520) (STJ110793)

STJ110793

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB:1:500-1:2000
Range	IHC-P:1:100-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	
Gene Symbol	RIPK4
Uniprot ID	RIPK4_HUMAN
Immunogen	
Immunogen Region	240-520
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 240-520 of human RIPK4 (NP_065690.2).
Immunogen Sequence	HRPELPVVCRRACSHLI RLMQRCWQGDPRVRPTFQEI TSETEDLCEKPDDEVKETAH DLDVKSPPPEPRSEVVPARLK RASAPTFDNDYSLSELLSQL DSGVQAVEGPEELSRSSSE SKLPSSGSGKRLSGVSSVDS AFSSRGLSLSFEREPTSD LGTTDVQKKLVDAIVSGDT SKLMKILQPQVDLALDSGA SLLHLAVEAGQECAKWLLL NNANPNLSNRRGSTPLHMA




Western blot analysis of extracts of A375 cells, using RIPK4 rabbit polyclonal antibody (STJ110793) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded human stomach using RIPK4 antibody (STJ110793) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry of paraffin-embedded human liver damage using RIPK4 antibody (STJ110793) at dilution of 1:100 (40x lens).



Immunohistochemistry analysis of paraffin-embedded human liver damage using RIPK4 antibody (STJ110793) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081