

## Anti-Phospho-ALK-Y1096 antibody (STJ110843)

STJ110843

### GENERAL INFORMATION


|                          |                    |
|--------------------------|--------------------|
| <b>Product Type</b>      | Primary antibodies |
| <b>Short Description</b> |                    |
| <b>Applications</b>      | WB/ELISA           |
| <b>Host/Source</b>       | Rabbit             |
| <b>Reactivity</b>        | Human              |

### PRODUCT PROPERTIES


|                            |   |
|----------------------------|---|
| <b>Clonality</b>           | Polyclonal  |
| <b>Clone ID</b>            |   |
| <b>Concentration</b>       | Lot specific  |
| <b>Conjugation</b>         | Unconjugated  |
| <b>Purification</b>        | Affinity purification   |
| <b>Dilution Range</b>      | WB:1:500-1:2000<br>ELISA:Recommended starting concentration is 1 $\mu$ g/mL. Please optimize the concentration based on your specific assay requirements. |
| <b>Formulation</b>         | PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.  |
| <b>Isotype</b>             | IgG   |
| <b>Storage Instruction</b> | Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.  |

### TARGET INFORMATION

|                           |   |
|---------------------------|---|
| <b>Gene ID</b>            | 238   |
| <b>Gene Symbol</b>        | ALK   |
| <b>Uniprot ID</b>         | ALK_HUMAN   |
| <b>Immunogen</b>          | PNYCF   |
| <b>Immunogen Region</b>   |   |
| <b>Specificity</b>        | A synthetic phosphorylated peptide around Y1096 of human ALK (NP_004295.2). |
| <b>Immunogen Sequence</b> | PNYCF   |



Western blot analysis of extracts of HL-60 and HL-60 cells, using Phospho-ALK-Y1096 antibody (STJ110843) at 1:1000 dilution. HL-60 cells were treated by 10% FBS after serum-starvation overnight. HL-60 cells were treated by UV for 15-30 minutes. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25 $\mu$ g per lane. Blocking buffer: 3% BSA.



Western blot analysis of lysates from HL-60 and HL-60 cells, using Phospho-ALK-Y1096 Rabbit polyclonal antibody (STJ110843) at 1:1000 dilution. HL-60 cells were treated by 10% FBS after serum-starvation overnight. HL-60 cells were treated by UV for 15-30 minutes. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25  $\mu$ g per lane. Blocking buffer: 3% BSA.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081