

**Anti-SERINC1 antibody (350-453) (STJ11100432)**  
STJ11100432

**GENERAL INFORMATION**


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	
<b>Applications</b>	WB/IHC-P/ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human/Mouse/Rat

**PRODUCT PROPERTIES**


<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	Lot specific
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purification
<b>Dilution Range</b>	WB:1:500-1:2000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
<b>Formulation</b>	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
<b>Isotype</b>	IgG
<b>Storage</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
<b>Instruction</b>	

**TARGET INFORMATION**


<b>Gene ID</b>	57515
<b>Gene Symbol</b>	SERINC1
<b>Uniprot ID</b>	SERC1_HUMAN
<b>Immunogen</b>	
<b>Immunogen Region</b>	350-453
<b>Specificity</b>	A synthetic peptide corresponding to a sequence within amino acids 350-453 of human SERINC1 (NP_065806.1).
<b>Immunogen Sequence</b>	ESTLIEDGGARSDGSLEDGD DVHRAVDNERDGVTSYSFF HFMLFLASLYIMMTLTNWYR YEPSREMKSQWTAVVVKISS SWGIVLYVWTLVAPLVLTN RDFF




Immunohistochemistry analysis of SERINC1 in paraffin-embedded Human liver using SERINC1 Rabbit polyclonal antibody (STJ11100432) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of SERINC1 in paraffin-embedded Mouse spleen using SERINC1 Rabbit polyclonal antibody (STJ11100432) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of SERINC1 in paraffin-embedded Rat spleen using SERINC1 Rabbit polyclonal antibody (STJ11100432) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of various lysates using SERINC1 Rabbit polyclonal antibody (STJ11100432) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBS-T. Detection: ECL Basic Kit. Exposure time: 10s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081