

Anti-Phospho-CTNNB1-S33/S37/T41 antibody (STJ11100967)

STJ11100967

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	IHC-P/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	1499
Gene Symbol	CTNNB1
Uniprot ID	CTNB1_HUMAN
Immunogen	DSGIHSGATT
Immunogen Region	
Specificity	A synthetic phosphorylated peptide around S33 & T41 of human beta Catenin (NP_001091679).
Immunogen Sequence	DSGIHSGATT




Western blot analysis of extracts of NIH/3T3 cells, using Phospho-Beta-Catenin-S33/S37/T41 antibody (STJ11100967) at 1:1000 dilution. NIH/3T3 cells were treated by Calyculin A (100 nM) at 37 °C for 30 minutes after serum-starvation overnight. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.



Immunohistochemistry analysis of Beta-Catenin in paraffin-embedded mouse kidney using Phospho-Beta-Catenin-S33/S37/T41 antibody (STJ11100967) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry of paraffin-embedded human colon carcinoma using Phospho-Beta-Catenin-S33/S37/T41 antibody (STJ11100967) at dilution of 1:100 (40x lens).



Immunohistochemistry analysis of Beta-Catenin in paraffin-embedded rat skin using Phospho-Beta-Catenin-S33/S37/T41 antibody (STJ11100967) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with immunohistochemistry staining protocol.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081