

Anti-ASNS antibody (450-550) [S7MR] (STJ11101267)

STJ11101267

GENERAL INFORMATION


Product Type Primary antibodies
Short Description
Applications WB/ELISA
Host/Source Rabbit
Reactivity Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality Monoclonal
Clone ID S7MR
Concentration Lot specific
Conjugation Unconjugated
Purification Affinity purification
Dilution Range WB:1:500-1:2000
ELISA: Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.
Isotype IgG
Storage Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
Instruction

TARGET INFORMATION

Gene ID 440
Gene Symbol ASNS
Uniprot ID ASNS_HUMAN
Immunogen
Immunogen 450-550
Region
Specificity A synthetic peptide corresponding to a sequence within amino acids 450-550 of human ASNS (P08243).
Immunogen ETFEDSNLIPKEILWRPKEA FSDGITSVKNSWFKILQEYV EHQVDDAMMANAAQKFPFNT PTKTEGYYYRQVFERHYPGR
Sequence ADWLSHYWMKPWINATDPSA R



Western blot analysis of extracts of MCF7 cells, using ASNS Rabbit monoclonal antibody (STJ11101267) at 1/141000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1/10000 dilution. Lanes/Proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 3min.



Western blot analysis of extracts of various cell lines, using ASNS Rabbit monoclonal antibody (STJ11101267) at 1/141000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1/10000 dilution. Lanes/Proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 3s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
 St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081