

Anti-ANXA2 antibody (1-100) [S0MR] (STJ11101300)

STJ11101300

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Monoclonal
Clone ID	S0MR
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:1000-1:6000 IHC-P:1:800-1:10000 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	302
Gene Symbol	ANXA2
Uniprot ID	ANXA2_HUMAN
Immunogen	
Immunogen Region	1-100
Specificity	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Annexin A2 (P07355).
Immunogen Sequence	MSTVHEILCKLSLEGDHSTP PSAYGSVKAYTNFDAERDAL NIETAIKTKGVDEVTIVNIL TNRSNAQRQDIAFAYQRRTK KELASALKSALSGHLETVIL




Western blot analysis of extracts of various cell lines, using Annexin A2 rabbit monoclonal antibody (STJ11101300) at 1:1000 dilution. Secondary antibody: HRP-Goat Anti-rabbit IgG (H+L) (STJ5000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.



Immunohistochemistry analysis of Annexin A2 in paraffin-embedded human cervix cancer tissue using Annexin A2 rabbit monoclonal antibody (STJ11101300) at a dilution of 1:800 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to immunohistochemistry staining.



Immunohistochemistry analysis of Annexin A2 in paraffin-embedded human colon carcinoma tissue using Annexin A2 rabbit monoclonal antibody (STJ11101300) at a dilution of 1:800 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to immunohistochemistry staining.



Immunohistochemistry analysis of Annexin A2 in paraffin-embedded mouse colon tissue using Annexin A2 rabbit monoclonal antibody (STJ11101300) at a dilution of 1:800 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to immunohistochemistry staining.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081