

**Anti-HLA-DQA1 antibody (155-254) [S5MR] (STJ11101305)**  
STJ11101305

**GENERAL INFORMATION**


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	
<b>Applications</b>	WB/IHC-P/ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human/Mouse/Rat

**PRODUCT PROPERTIES**


<b>Clonality</b>	Monoclonal
<b>Clone ID</b>	S5MR
<b>Concentration</b>	Lot specific
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purification
<b>Dilution Range</b>	WB:1:1000-1:6000 IHC-P:1:200-1:2000 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
<b>Formulation</b>	PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.
<b>Isotype</b>	IgG
<b>Storage Instruction</b>	Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

**TARGET INFORMATION**


<b>Gene ID</b>	3117
<b>Gene Symbol</b>	HLA-DQA1
<b>Uniprot ID</b>	DQA1_HUMAN
<b>Immunogen</b>	
<b>Immunogen Region</b>	155-254
<b>Specificity</b>	A synthetic peptide corresponding to a sequence within amino acids 155-254 of human HLA-DQA1 (P01909). EGVSETSFSLKSDHSFFKIS YLTLLPSAEESYDCKVEHWG LDKPLLKHWEPEIPAPMSEL TETVVCALGLSVGLVGIIVGG
<b>Immunogen Sequence</b>	TVFIIRGLRSVGASRHQGPL




Western blot analysis of extracts of various cell lines, using HLA-DQA1 rabbit monoclonal antibody (STJ11101305) at 1:1000 dilution. Secondary antibody: HRP-Goat Anti-rabbit IgG (H+L) (STJ5000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.



Immunohistochemistry analysis of HLA-DQA1 in paraffin-embedded human colon tissue using HLA-DQA1 rabbit monoclonal antibody (STJ11101305) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to immunohistochemistry staining.



Immunohistochemistry analysis of HLA-DQA1 in paraffin-embedded human liver tissue using HLA-DQA1 rabbit monoclonal antibody (STJ11101305) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to immunohistochemistry staining.



Immunohistochemistry analysis of HLA-DQA1 in paraffin-embedded human tonsil tissue using HLA-DQA1 rabbit monoclonal antibody (STJ11101305) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to immunohistochemistry staining.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081