

## Anti-SUPT16H antibody (520-695) (STJ11101487) STJ11101487

## **GENERAL INFORMATION**

Primary antibodies
WB/IF/ICC/ELISA
Rabbit
Human/Mouse/Rat

## **PRODUCT PROPERTIES**

Polyclonal
Lot specific
Unconjugated
Affinity purification
WB:1:500-1:1000
IF/ICC:1:50-1:200
ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
IgG
Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.


## **TARGET INFORMATION**

Gene ID 11198 Gene Symbol SUPT16H Immunogen Immunogen 520-695 Region

Uniprot ID SP16H\_HUMAN


Specificity Recombinant fusion protein containing a sequence corresponding to amino acids 520-695 of human SUPT16H (SPT16) (NP\_009123.1).

Immunogen KEPHIREMKIYIDKKYETVI MPVFGIATPFHIATIKNISM SVEGDYTYLRINFYCPGSAL GRNEGNIFPNPEATFVKEIT Sequence YRASNIKAPGEQTVPALNLQ NAFRIIKEVQKRYKTREAEE KEKEGIVKQDSLVINLNRSN PKLKDLYIRPNIAQKRMQGS LEAHVNGFRFTSVRGD



blot analysis of various lysates, u (SPT16) rabbit polyclonal antii (487) at 1:400 dilution. Secondary antib at Anti-rabbit IgG (H+L) (STJS000856 n. Lysa : 3% no arproteins: 25 Mu g per fat dry milk in TBST. Dete time: 60s.

Immunofluorescence analysis of U2OS using SUPT16H (SPT16) antibody (STJ11101487) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 using SUPT16H (SPT16) antibody (STJ11101487) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes. St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081