

Anti-TSC2 antibody (1700-1807) [S7MR] (STJ11101687)

STJ11101687

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications WB/IP/ELISA
Host/Source Rabbit
Reactivity Human

PRODUCT PROPERTIES

Clonality Monoclonal S7MR
Concentration Lot specific Unconjugated Purification
Purification
Dilution Range WB:1:500-1:1000

IP:0.5 Mu g-4 Mu g antibody for 200 Mu g-400 Mu g extracts of whole cells

 $ELISA: Recommended \ starting \ concentration \ is \ 1 \ Mu \ g/mL. \ Please \ optimize \ the \ concentration \ based \ on \ your \ specific \ assay$

requirements.

Formulation PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.

Isotype IgG

Storage Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

Instruction

TARGET INFORMATION

Gene ID 7249
Gene Symbol TSC2
Uniprot ID TSC2_HUMAN


Immunogen

Immunogen 1700-1807 Region


Specificity A synthetic peptide corresponding to a sequence within amino acids 1700-1807 of human Tuberin (P49815).

Immunogen AKIVSDRNLPFVARQMALHA NMASQVHHSRSNPTDIYPSK WIARLRHIKRLRQRICEEAA YSNPSLPLVHPPSHSKAPAQ

Sequence TPAEPTPGYEVGQRKRLISS VEDFTEFV



Immunoprecipitation analysis of 300 Mu g extracts of 293T cells using 3 Mu g TSC2 antibody (STJ11101687). Western blot was performed from the immunoprecipitate using TSC2 antibody (STJ11101687) at a dilution of 1:1000.



Western blot analysis of various lysates using TSC. Rabbit monoclonal antibody (STJ.11101687) at 1:1001 dilution. Secondary antibody: HRP Goat Anti-Rabbi IgG (H+L) (STJS000856) at 1:10000 dilution Lysates/proteins: 25 Mu g per lane. Blocking buffer: 39 nonfat dry milk in TBST. Detection: ECL Basic Kit Exposure fine: 016