

Anti-ACVR1B antibody (1-100) [S4MR] (STJ11101804)

ST.111101804

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications WB/IHC-P/IF/ICC/ELISA

Host/Source Rabbit

Reactivity Human/Mouse/Rat

PRODUCT PROPERTIES

Clonality Clone ID
Monoclonal S4MR

Concentration Conjugation Purification
Lot specific Unconjugated Principle Pri

IF/ICC:1:200-1:400

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.

Isotype IgG

Storage Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

Instruction


TARGET INFORMATION

Immunogen 1-100 Region


Specificity A synthetic peptide corresponding to a sequence within amino acids 1-100 of human ACVR1B (P36896).

Immunogen MAESAGASSFFPLVVLLLAG SGGSGPRGVQALLCACTSCL QANYTCETDGACMVSIFNLD GMEHHVRTCIPKVELVPAGK


Sequence PFYCLSSEDLRNTHCCYTDY



Western blot analysis of extracts of various cell lines, using ACVR1B rabbit monoclonal antibody (STJ11101804) at 1:1000 dilution. Secondary antibody HRP Goat Artit-rabbit IgG (H+L) (STJS00056) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection



Immunohistochemistry, analysis of paraffin-embedded rat testis using ACVR1B rabbit monocional antibody (STJ11101804) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM fris/EDTA buffer pH 9. 0 before commencing with



Immunohistochemistry analysis of paraffin-embedder human liver using ACVP1B rabbit monoclonal antibody (STJ11101804) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9. 0 before commencing with



Immunohistochemistry analysis of paraffin-embedded mouse brain using ACVR1B rabbit monocional antibody (STJ11101804) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9. 0 before commencing with