

## Anti-SNCG antibody (48-127) [S1MR] (STJ11101831)

ST.I1110183

## **GENERAL INFORMATION**

Product Type Primary antibodies

**Short Description** 

Applications WB/IHC-P/ELISA Host/Source Rabbit

Reactivity Human/Mouse/Rat

## **PRODUCT PROPERTIES**

Clonality Monoclonal
Clone ID S1MR

Concentration Lot specific
Conjugation Unconjugated
Purification Affinity purification
Dilution Range WB:1:500-1:1000

IHC-P:1:50-1:200

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.

**Isotype** IgG

**Storage Instruction** Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

## **TARGET INFORMATION**


Gene ID 6623
Gene Symbol SNCG
Uniprot ID SYUG\_HUMAN
Immunogen

Immunogen Region 48-127


Specificity A synthetic peptide corresponding to a sequence within amino acids 48-127 of human gamma-Synuclein (O76070).

Immunogen Sequence

VVQSVTSVAEKTKEQANAVS EAVVSSVNTVATKTVEEAEN IAVTSGVVRKEDLRPSAPQQ EGEASKEKEEVAEEAQSGGD Sequence



Immunohistochemistry analysis of paraffin-embedden rat brain using gamma-Synuclein Rabbit monoclona antibody (STJ11101831) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with



Western blot analysis of extracts of various cell lines using gamma-Synuclein Rabbit monoclonal antibody (STJ11101831) at 11%1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H-L (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk ir TBST Datesfore SCI Rapie Kiff Exposure time; Smir