

## Anti-RAP1GAP antibody (1-100) [S2MR] (STJ11101912)

ST.I11101912

## **GENERAL INFORMATION**

Product Type Primary antibodies

**Short Description** 

Applications WB/IF/ICC/ELISA Host/Source Rabbit

Reactivity Human/Mouse/Rat

## **PRODUCT PROPERTIES**

Clonality Monoclonal S2MR

Concentration Lot specific Unconjugated Purification Pulling WB:1:500-1:1000

IF/ICC:1:50-1:200

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.

**Isotype** IgG

**Storage Instruction** Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

## **TARGET INFORMATION**

Gene ID 5909

Gene Symbol RAP1GAP Uniprot ID RPGP1\_HUMAN


. Immunogen

Immunogen Region 1-100


Specificity A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Rap1GAP (P47736).

 ${\bf Immunogen} \quad {\tt MIEKMQGSRMDEQRCSFPPP\,LKTEEDYIPYPSVHEVLGRE\,GPFPLILLPQFGGYWIEGTN\,HEITSIPETEPLQSPTTKVK}$ 

Sequence LECNPTARIYRKHFLGKEHF



Immunofluorescence analysis of NIH-3T3 cells using Rap1GAP Rabbit monoclonal antibody (STJ11101912 at dilution of 1:100 (40x lens). Blue: DAPI for nuclea staining.



Western blot analysis of extracts of various cell lines, using Rap1GAP Rabbit monoclonal artibody (STJ11101912) at 1:1000 (dilution. Secondary antibody; HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% norflat dry milk in TBST. Detection: