

Anti-NDRG1 antibody (295-394) [S6MR] (STJ11101986)
STJ11101986

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Monoclonal
Clone ID	S6MR
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	10397
Gene Symbol	NDRG1
Uniprot ID	NDRG1_HUMAN
Immunogen	
Immunogen Region	295-394
Specificity	A synthetic peptide corresponding to a sequence within amino acids 295-394 of human NDRG1 (Q92597).
Immunogen Sequence	ISQPAKLAEAFKYFVQGMGY MPSASMTLRMSRTASGSSV TSLDGTRSRSHSTSEGTRSRSH TSEGTRSRSHSTSEGAHLDI TPNSGAAGNSAGPKSMEVSC




Western blot analysis of extracts of various cell lines, using NDRG1 rabbit monoclonal antibody (STJ11101986) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.



Immunohistochemistry analysis of NDRG1 in paraffin-embedded human colon carcinoma tissue using NDRG1 rabbit monoclonal antibody (STJ11101986) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to immunohistochemistry staining.



Immunohistochemistry analysis of NDRG1 in paraffin-embedded human lung squamous carcinoma tissue using NDRG1 rabbit monoclonal antibody (STJ11101986) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to immunohistochemistry staining.



Immunohistochemistry analysis of NDRG1 in paraffin-embedded human tonsil tissue using NDRG1 rabbit monoclonal antibody (STJ11101986) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to immunohistochemistry staining.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081