

Anti-CCNH antibody (224-323) [S1MR] (STJ11101991)

STJ11101991

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse

PRODUCT PROPERTIES


Clonality	Monoclonal
Clone ID	S1MR
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	902
Gene Symbol	CCNH
Uniprot ID	CCNH_HUMAN
Immunogen	
Immunogen Region	224-323
Specificity	A synthetic peptide corresponding to a sequence within amino acids 224-323 of human Cyclin H (P51946).
Immunogen Sequence	AGITMESYLSLMLKENRT CLSQLDIMKSMRNLVKYK PPRSEEVAVLKQKLERCHSA ELALNVITKKRKGYEDDDYV SKKSKHEEEWTDLDLVEL




Immunofluorescence analysis of U-2 OS cells using Cyclin H Rabbit monoclonal antibody (STJ11101991) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH-3T3 cells using Cyclin H Rabbit monoclonal antibody (STJ11101991) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded mouse testis using Cyclin H Rabbit monoclonal antibody (STJ11101991) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using Cyclin H Rabbit monoclonal antibody (STJ11101991) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 60s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081