

Anti-CCT2 antibody (436-535) [S4MR] (STJ11102134)

STJ11102134

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications
Host/Source
Rabbit
Human/Mouse/Rat

PRODUCT PROPERTIES

Clonality Monoclonal
Clone ID S4MR

Concentration Lot specific
Conjugation Unconjugated
Purification Affinity purification
Dilution Range WB:1:500-1:2000

IHC-P:1:50-1:200

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.

Isotype IgG

Storage Instruction Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID 10576
Gene Symbol CCT2
Uniprot ID TCPB_HUMAN
Immunogen

Immunogen Region 436-535


Specificity A synthetic peptide corresponding to a sequence within amino acids 436-535 of human TCP1 beta/CCT2 (P78371).

Immunogen MESYAKALRMLPTIIADNAG YDSADLVAQLRAAHSEGNTT AGLDMREGTIGDMAILGITE SFQVKRQVLLSAAEAAEVIL

Sequence RVDNIIKAAPRKRVPDHHPC



Immunohistochemistry analysis of paraffin-embedder rat brain using TCP1 beta/CCT2 Rabbit monoclona antibody (STJ11102134) at dilution of 1:100 (40x lens) Perform microwave antigen retireval with 10 mM PBS buffer pH 7. 2 before commencing with



Western blot analysis of extracts of various cell lines, sing TCP1 beta/CCT2 Rabbit monoclonal antibody; STJ11102134] at 1:1000 dilution. Secondary antibody; HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: 2CL Basic Ktt. Exposure time: 3s.