

Anti-MSN antibody (400-500) [S6MR] (STJ11102196)

STJ11102196

GENERAL INFORMATION


| | |
|--------------------------|--------------------|
| Product Type | Primary antibodies |
| Short Description | |
| Applications | WB/IHC-P/ELISA |
| Host/Source | Rabbit |
| Reactivity | Human/Mouse/Rat |

PRODUCT PROPERTIES


| | |
|----------------------------|--|
| Clonality | Monoclonal |
| Clone ID | S6MR |
| Concentration | Lot specific |
| Conjugation | Unconjugated |
| Purification | Affinity purification |
| Dilution Range | WB:1:500-1:2000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements. |
| Formulation | PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3. |
| Isotype | IgG |
| Storage Instruction | Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles. |

TARGET INFORMATION


| | |
|---------------------------|---|
| Gene ID | 4478 |
| Gene Symbol | MSN |
| Uniprot ID | MOES_HUMAN |
| Immunogen | |
| Immunogen Region | 400-500 |
| Specificity | A synthetic peptide corresponding to a sequence within amino acids 400-500 of human MSN (P26038). |
| Immunogen Sequence | KEALLQASRDQKKTQEQLAL EMAELTARISQLEMARQKKE SEAVEWQKQAMVQEDLEKT RAEKLTAMSTPHVAEPAENE QDEQDENGAEASADLRADAM A |



Immunohistochemistry analysis of paraffin-embedded human colon using Moesin Rabbit monoclonal antibody (STJ11102196) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat spleen using Moesin Rabbit monoclonal antibody (STJ11102196) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using Moesin Rabbit monoclonal antibody (STJ11102196) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081