

Anti-ADA antibody (250-350) [S6MR] (STJ11102276)

ST.J11102276

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications WB/ELISA Host/Source Rabbit

Reactivity Human/Mouse/Rat

PRODUCT PROPERTIES

Clonality Monoclonal
Clone ID S6MR

Concentration Lot specific
Conjugation Unconjugated
Purification Affinity purification
Dilution Range WB:1:500-1:1000

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.

Isotype IgG

Storage Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

Instruction

TARGET INFORMATION

Gene ID 100 Gene Symbol ADA

Uniprot ID ADA_HUMAN

Immunogen

Immunogen 250-350


Region

Specificity A synthetic peptide corresponding to a sequence within amino acids 250-350 of human Adenosine Deaminase (ADA)


(NP 000013.2)

Immunogen NRLRQENMHFEICPWSSYLT GAWKPDTEHAVIRLKNDQAN YSLNTDDPLIFKSTLDTDYQ MTKRDMGFTEEEFKRLNINA

Sequence AKSSFLPEDEKRELLDLLYK A



Western blot analysis of various lysates, using Adenosine Deaminase (ADA) rabbit monoclonal antibody (STJ11102276) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mug per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection. ECC Resic Kit Exponers into 6the



Western blot analysis of mouse stomach, using Adenosine Deaminase (ADA) rabbit monoclonal antibody (STJ11102276) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mug per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection ECL Enbanced Mr. Exposure time: 45c.