

**Anti-hnRNP C antibody (80-170) [S2MR] (STJ11102402)**  
STJ11102402

**GENERAL INFORMATION**


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	
<b>Applications</b>	WB/IF/ICC/ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human/Mouse

**PRODUCT PROPERTIES**

<b>Clonality</b>	Monoclonal
<b>Clone ID</b>	S2MR
<b>Concentration</b>	Lot specific
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purification
<b>Dilution Range</b>	WB:1:1000-1:6000 IF/ICC:1:200-1:2000 ELISA:Recommended starting concentration is 1 $\mu$ g/mL. Please optimize the concentration based on your specific assay requirements.
<b>Formulation</b>	PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.
<b>Isotype</b>	IgG
<b>Storage Instruction</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.


**TARGET INFORMATION**

<b>Gene ID</b>	3183
<b>Gene Symbol</b>	HNRPNC
<b>Uniprot ID</b>	HNRPNC_HUMAN
<b>Immunogen</b>	
<b>Immunogen Region</b>	80-170
<b>Specificity</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 80-170 of human hnRNP C (P07910). LDINLAAEPKVNRGKAGVKR SAAEMYGSVTEHPSPSPLLS SSFDLDYDFQRDYYDRMYSY PARVPPPPPIARAVVPSKRQ
<b>Sequence</b>	RVSGNTSRRGK




Western blot analysis of extracts of various cell lines, using hnRNP C rabbit monoclonal antibody (STJ11102402) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25  $\mu$ g per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 3s.

Confocal imaging of a paraffin-embedded Human colon cancer using hnRNP C Rabbit monoclonal antibody (STJ11102402, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 40x. Perform high pressure antigen retrieval with 0.01 M citrate buffer (pH 6.0) prior to IF staining.



Immunofluorescence analysis of U-2 OS cells using hnRNP C rabbit monoclonal antibody (STJ11102402) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Confocal imaging of U-2 OS cells using hnRNP C Rabbit monoclonal antibody (STJ11102402, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (dilution 1:500) (Red). The cells were counterstained with Alpha-Tubulin Mouse monoclonal antibody (dilution 1:400) followed by incubation with ABFluor 488-conjugated Goat Anti-Mouse IgG (H+L) antibody (dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081