

Anti-ANXA5 antibody (221-320) [S5MR] (STJ11102465)

ST.111102465

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications WB/ELISA Host/Source Rabbit

Reactivity Human/Mouse/Rat

PRODUCT PROPERTIES

Clonality Monoclonal
Clone ID S5MR

Concentration Lot specific
Conjugation Unconjugated
Purification Affinity purification
Dilution Range WB:1:500-1:2000

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements

Formulation PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.

Isotype IgG

Storage Instruction Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

 Gene ID
 308

 Gene Symbol
 ANXA5

 Uniprot ID
 ANXA5_HUMAN


 Immunogen
 221-320

Specificity A synthetic peptide corresponding to a sequence within amino acids 221-320 of human Annexin A5/Annexin A5/Annexin V


IP_001145.1)

Immunogen IEETIDRETSGNLEQLLLAV VKSIRSIPAYLAETLYYAMK GAGTDDHTLIRVMVSRSEID LFNIRKEFRKNFATSLYSMI


Sequence KGDTSGDYKKALLLLCGEDD




Western blot analysis of extracts of various cell lines, using Annexin A5/Annexin A5/Annexin V rabbit monoclonal antibody (STJ1102465) at 1:1000 dilution Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection



Western blot analysis of extracts of various cell lines, using Annexin A5/Annexin A5/Annexin V Rabbit monoclonal antibody (St11102465) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in



Western blot analysis of extracts of various cell lines, using Annexin A5/Annexin A5/Annexin V rabbit monoclonal antibody (STJ1102465) at 1:1000 dilution Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane Blocking buffer: 3% non-lat dry milk in TBST. Detection



Western blot analysis of extracts of various cell lines using Annexin A5/Annexin A5/Annexin V Raio monoclonal antibody (STJ11102/465) at 1:100 Raibo Secondary antibody: HRP Goat Anti-Raibbit IgG (H-ti (STJS000856) at 1:10000 dilution. Lysates/proteins: 2! Mu g per lane. Blocking buffer: 3% nonfat dry milk in ITST. Detection: ECL Basic Kit. Exoosure time: 1s.