

## Anti-TRMT2A/HTF9C antibody (1-100) [S4MR] (STJ11102674)

ST.I11102674

## **GENERAL INFORMATION**

Product Type Primary antibodies

**Short Description** 

Applications WB/ELISA
Host/Source Rabbit
Reactivity Human

## **PRODUCT PROPERTIES**

Clonality Monoclonal S4MR
Concentration Lot specific Unconjugated Purification Pullution Range WB:1:500-1:1000

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.

**Isotype** IgG

Storage Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

Instruction

## **TARGET INFORMATION**


Gene ID 27037
Gene Symbol TRMT2A
Uniprot ID TRM2A\_HUMAN

Immunogen Immunogen 1-100 Region


Specificity A synthetic peptide corresponding to a sequence within amino acids 1-100 of human TRMT2A/HTF9C (Q8IZ69).

Immunogen MSENLDNEGPKPMESCGQES SSALSCPTVSVPPAAPAALE EVEKEGAGAATGPGPQPGLY SYIRDDLFTSEIFKLELQNV

Sequence PRHASFSDVRRFLGRFGLQP



Western blot analysis of K-562, using TRMT2A/HTF9(rabbit monoclonal antibody (STJ11102674) at 1:100 dilution. Secondary antibody: HRP Goat Anti-rabbit (gl-(H+L) (STJS000856) at 1:10000 dilutior. Lysates/proteins: 25ug per lane. Blocking buffer: 39 non-fat dry milk in TBST. Detection: ECL Basic Kif



Western blot analysis of MCF7, using TRMT2A/HTF9C rabbit monoclonal antibody (ST11102674) at 1:1000 (dilution. Secondary antibody: HHP Goat Anti-rabbit (gG H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% non-fat dry milk in 'HST. Detection: ECL Basic Kit.