

Anti-ACAA2 antibody (298-397) [S4MR] (STJ11102704)

STJ11102704

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications WB/IF/ICC/ELISA Host/Source Rabbit

Reactivity Human/Mouse/Rat

PRODUCT PROPERTIES

Clonality Monoclonal
Clone ID S4MR

Concentration Lot specific
Conjugation Unconjugated
Purification Affinity purification
Dilution Range WB:1:500-1:1000

IF/ICC:1:50-1:200

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.

Isotype IgG

Storage Instruction Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID 10449
Gene Symbol ACAA2
Uniprot ID THIM_HUMAN
Immunogen

Immunogen Region 298-397


Specificity A synthetic peptide corresponding to a sequence within amino acids 298-397 of human ACAA2 (P42765).

Immunogen PAISGALKKAGLSLKDMDLV EVNEAFAPQYLAVERSLDLD ISKTNVNGGAIALGHPLGGS GSRITAHLVHELRRRGGKYA


Sequence VGSACIGGGQGIAVIIQSTA




Immunofluorescence analysis of U2OS cells using ACAA2 Rabbit monoclonal antibody (STJ11102704) al dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells usin ACAA2 Rabbit monoclonal antibody (STJ11102704) a dilution of 1:100 (40x lens). Blue: DAPI for nuclear statistics



Immunofluorescence analysis of HepG2 cells using ACAA2 Rabbit monoclonal antibody (STJ11102704) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear extension



Western blot analysis of extracts of various cell lines using ACA2 antibody (STJ11102704) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbi IgG (H+L) (STJS000856) at 1:10000 dilution Lysates/proteins: 25 Mu g per lane. Blocking buffer 3% nonfat dry rnilk in TBST. Detection: ECL Basic Kit