

Anti-UBE4A antibody (1-100) [S7MR] (STJ11102707)

STJ11102707

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications WB/ELISA Host/Source Rabbit

Reactivity Human/Mouse/Rat

PRODUCT PROPERTIES

Clonality Monoclonal
Clone ID S7MR

Concentration Lot specific
Conjugation Unconjugated
Purification Affinity purification
Dilution Range WB:1:500-1:1000

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.

Isotype IgG

Storage Instruction Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID 9354 Gene Symbol UBE4A

ene Symbol UBE4A Uniprot ID UBE4A_HUMAN


Immunogen

Immunogen Region 1-100


Specificity A synthetic peptide corresponding to a sequence within amino acids 1-100 of human UBE4A (Q14139).

Immunogen MTDQENNNNISSNPFAALFG SLADAKQFAAIQKEQLKQQS DELPASPDDSDNSVSESLDE FDYSVAEISRSFRSQQEICE

Sequence QLNINHMIQRIFLITLDNSD



Western blot analysis of extracts of various cell line using UBE4A antibody (STJ11102707) at 1:100 dilution. Secondary antibody: HRP Goat Anti-Rabl IgG (H+L) (STJS000856) at 1:10000 dilutic Lysates/profiers: 25 Mu g per lane. Blocking buffer 3 nontat dry milk in TBST. Detection: ECL Basic K



Western blot analysis of extracts of HUVEC cells, using UBE4A antibody (STJ11102707) at 1:1000 dilution Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L (STJS000856) at 1:10000 dilution. Lysates/proteins 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time; 900.