

Anti-Aromatase antibody (1-100) [S4MR] (STJ11102744)

STJ11102744

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications WB/IHC-P/IF/ICC/ELISA

Host/Source Rabbit Reactivity Human

PRODUCT PROPERTIES

IHC-P:1:800-1:8000 IF/ICC:1:100-1:400

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.

Isotype Ig0

Storage Instruction Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID 1588

Gene Symbol CYP19A1
Uniprot ID CP19A_HUMAN

Immunogen


gen

Immunogen Region 1-100


Specificity A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Aromatase (CYP19A1) (P11511).

Immunogen MVLEMLNPIHYNITSIVPEA MPAATMPVLLLTGLFLLVWN YEGTSSIPGPGYCMGIGPLI SHGRFLWMGIGSACNYYNRV


Sequence YGEFMRVWISGEETLIISKS



Immunofluorescence analysis of human placenta using Aromatase (CYP19A1) Rabbit monoclonal antibody (STJ11102744) at dilution of 1:100 (40x lens). Blue DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded duman colon using Aromatase (CYP19A1) Rabbit monoclonal antibody (STJ11102744) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with



Western blot analysis of extracts of U-87MG cells using Aromatase (CYP19A1) antibody (STJ11102744) at 1:1000 dilution. Secondary antibody: HIPR Goat Anti Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution Lysates/profesirs: 25 Mu g per lane. Blocking buffer: 39 nonfat dry milk in TBST. Detection: ECL Enhanced Kit Evonsure time: 180s.