

Anti-PDGFRB antibody (900-1000) (STJ11102763)

STJ11102763

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	5159
Gene Symbol	PDGFRB
Uniprot ID	PGFRB_HUMAN
Immunogen	
Immunogen Region	900-1000
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 900-1000 of human PDGFRB (NP_002600.1).
Immunogen Sequence	GGTPYPELPMNEQFYNAIKR GYRMAQPAHASDEIYEIMQK CWEEKFEIRPPFSQLVLLLE RLLGEGYKKKYQQVDEEFLR SDHPAILRSQARLPGFHGLR S




Immunofluorescence analysis of NIH-3T3 cells using PDGFRB Rabbit polyclonal antibody (STJ11102763) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of PDGFRB in paraffin-embedded human esophageal cancer using PDGFRB Rabbit polyclonal antibody (STJ11102763) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6, 0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of PDGFRB in paraffin-embedded human lung cancer using PDGFRB Rabbit polyclonal antibody (STJ11102763) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6, 0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of lysates from SH-SY5Y cells, using PDGFRB Rabbit polyclonal antibody (STJ11102763) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ5000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 180s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081