

Anti-PDCD7 antibody (1-100) [S8MR] (STJ11102768)

ST.111102768

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications WB/IHC-P/ELISA
Host/Source Rabbit
Reactivity Human/Mouse/Rat

PRODUCT PROPERTIES

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

 $\begin{tabular}{ll} \hline \textbf{Formulation} & \dot{\text{PBS}} \text{ with } 0.02\% \text{ Sodium Azide, } 0.05\% \text{ BSA, } 50\% \text{ Glycerol, pH } 7.3. \\ \hline \end{tabular}$

Isotype IgG

Storage Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

Storage Instruction

TARGET INFORMATION

Gene ID 10081 Gene Symbol PDCD7


Uniprot ID PDCD7_HUMAN

Immunogen Immunogen 1-100 Region


Specificity A synthetic peptide corresponding to a sequence within amino acids 1-100 of human PDCD7 (Q8N8D1).

Immunogen MALPPFFGQGRPGPPPPQPP PPAPFGCPPPPLPSPAFPPP LPQRPGPFPGASAPFLQPPL ALQPRASAEASRGGGGAGAF

Sequence YPVPPPPLPPPPPQCRPFPG



Immunohistochemistry analysis of A PDCD7 in parafit embedded4 rat liver tissue using A PDCD7 Rabt monoclonal antibody (STJ11102768) at a ciliution 1:200 (40x lens). High pressure antigen retrieval w performed with 0, 01 M citrate buffer (pH 6. 0) prior immunohistochemistry training.



Immunohistochemistry analysis of PDCD7 in paraffic embeddedÅ human esophagus tissue usingÅ PDCD Rabbit monoclonal antibody (STJ11102768) at dilution of 1:200 (40x lens). High pressure antige retrieval was performed with 0. 01 M citrate buffer (p 6. 0) prior to immunohistochemistry staining.



mmunohistochemistry analysis ofÅ PDCD7 in paraffin ambeddedÅ mouse intestin tissue usingÅ PDCD. Rabbit monoclonal antibody (STJ11102768) at: dilution of 1:200 (40x lens). High pressure antiger etrieval was performed with 0.01 M citrate buffer (ph. 5.0 prior to immunohistochemistry staining.



Immunohistochemistry analysis of Å PDCD7 in paraffinembeddedÅ human small intestine tissue usingÅ PDCD7 Rabbit monoclonal antibody (STJ11102768) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to immunohistochemistry stainino.