

Anti-WFDC1 antibody (111-220) (STJ11102875)

STJ11102875

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	58189
Gene Symbol	WFDC1
Uniprot ID	WFDC1_HUMAN
Immunogen	
Immunogen Region	111-220
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 111-220 of human WFDC1 (NP_067020.2).
Immunogen Sequence	PPVLDWLQPKPRWLGGNGW LLDGPEEVLQAEACSTTDEG AEPLLCPSGYECHILSPGDV AEGIPNRGQCVKQRRQADGR ILRHKLYKEYPEGDSKNVAE PGRGQKHFQ



Immunofluorescence analysis of L929 cells using WFDC1 Rabbit polyclonal antibody (STJ11102875) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using WFDC1 Rabbit polyclonal antibody (STJ11102875) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.