

## Anti-PRKCDBP antibody (75-160) (STJ11102892)

STJ11102892

### GENERAL INFORMATION


|                          |                    |
|--------------------------|--------------------|
| <b>Product Type</b>      | Primary antibodies |
| <b>Short Description</b> |                    |
| <b>Applications</b>      | WB/ELISA           |
| <b>Host/Source</b>       | Rabbit             |
| <b>Reactivity</b>        | Rat                |

### PRODUCT PROPERTIES

|                            |   |
|----------------------------|---|
| <b>Clonality</b>           | Polyclonal  |
| <b>Clone ID</b>            |   |
| <b>Concentration</b>       | Lot specific  |
| <b>Conjugation</b>         | Unconjugated  |
| <b>Purification</b>        | Affinity purification   |
| <b>Dilution Range</b>      | WB:1:500-1:2000<br>ELISA:Recommended starting concentration is 1 $\mu$ g/mL. Please optimize the concentration based on your specific assay requirements. |
| <b>Formulation</b>         | PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.  |
| <b>Isotype</b>             | IgG   |
| <b>Storage Instruction</b> | Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.  |

### TARGET INFORMATION

|                           |  |
|---------------------------|--|
| <b>Gene ID</b>            | 112464   |
| <b>Gene Symbol</b>        | CAVIN3   |
| <b>Uniprot ID</b>         | CAVN3_HUMAN  |
| <b>Immunogen</b>          |  |
| <b>Immunogen Region</b>   | 75-160   |
| <b>Specificity</b>        | Recombinant fusion protein containing a sequence corresponding to amino acids 75-160 of human PRKCDBP (NP_659477.2). |
| <b>Immunogen Sequence</b> | SNTLAQLLAKAERVSSHANA AQERAVRRAAQVQRLEANHG LLVARGKLHVLLFKEEGEVP ASAFQKAPEPLGPADQSELG PEQLEA                           |



Western blot analysis of lysates from Rat heart, using PRKCDBP Rabbit polyclonal antibody (STJ11102892) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25  $\mu$ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081