

Anti-LPP antibody (1-100) [S8MR] (STJ11102958)

STJ11102958

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications WB/IHC-P/IF/ICC/ELISA

Host/Source Rabbit

Reactivity Human/Mouse/Rat

PRODUCT PROPERTIES

IF/ICC:1:50-1:200

 $ELISA: Recommended \ starting \ concentration \ is \ 1 \ Mu \ g/mL. \ Please \ optimize \ the \ concentration \ based \ on \ your \ specific \ assay$

requirements.

Formulation PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.

Isotype IgG

Storage Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

Instruction

TARGET INFORMATION

Gene ID 4026
Gene Symbol LPP
Uniprot ID LPP_HUMAN

Immunogen Immunogen 1-100 Region


Specificity A synthetic peptide corresponding to a sequence within amino acids 1-100 of human LPP (Q93052).

Immunogen MSHPSWLPPKSTGEPLGHVP ARMETTHSFGNPSISVSTQQ PPKKFAPVVAPKPKYNPYKQ PGGEGDFLPPPPPPLDDSSA


Sequence LPSISGNFPPPPPLDEEAFK



Immunofluorescence analysis of NIH/3T3 cells using LPP Rabbit monoclonal antibody (STJ11102958) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded human liver cancer using LPP Rabbit monoclona antibody (STJ11102958) at dilution of 1:100 (40x lens) Perform microwave antigen retrieval with 10 mlv Tris/EDTA buffer pH 9. 0 before commencing with



Western blot analysis of extracts of various cell lines, using LPP antibody (STJ11102958) at 1:1000 dillution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dillution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit Exposure time: 5s.