

Anti-HNRNPA3 antibody (100-200) (STJ11103228)
STJ11103228

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	220988
Gene Symbol	HNRNPA3
Uniprot ID	ROA3_HUMAN
Immunogen	
Immunogen Region	100-200
Specificity	A synthetic peptide corresponding to a sequence within amino acids 100-200 of human HNRNPA3 (NP_001317178.1).
Immunogen Sequence	VDGRVVEPKRAVSREDSVKP GAHLTVKKIFVGGIKEDTEE YNLRDYFEKYGIETIEVME DRQSGKKRGFAFVTFDDHDT VDKIVQKYHTINGHNCEVK K




Immunohistochemistry analysis of HNRNPA3 in paraffin-embedded mouse brain using HNRNPA3 Rabbit polyclonal antibody (STJ11103228) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of HNRNPA3 in paraffin-embedded human lung cancer using HNRNPA3 Rabbit polyclonal antibody (STJ11103228) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of HNRNPA3 in paraffin-embedded rat ovary using HNRNPA3 Rabbit polyclonal antibody (STJ11103228) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of lysates from U-251MG cells, using HNRNPA3 Rabbit polyclonal antibody (STJ11103228) at 1:1000 dilution. Secondary antibody: HRP-Goat Anti-Rabbit IgG (H+L) (STJ5000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081