

Anti-67kDa Laminin Receptor antibody (196-295) [S0MR] (STJ11103600)

STJ11103600

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Monoclonal
Clone ID	S0MR
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
Instruction	

TARGET INFORMATION


Gene ID	3921
Gene Symbol	RPSA
Uniprot ID	RSSA_HUMAN
Immunogen	
Immunogen	196-295
Region	
Specificity	A synthetic peptide corresponding to a sequence within amino acids 196-295 of human Laminin Receptor (P08865).
Immunogen Sequence	EVMPDLYFYRDPPEIEKEEQ AAAEKAVTKEEFQGEWTAPA PEFTATQPEVADWSEGVQVP SVPIQQFPTEDWSAQPATED



Confocal imaging of HeLa cells using 67kDa Laminin Receptor Rabbit monoclonal antibody (STJ11103600, dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (dilution 1:500) (Red). The cells were counterstained with Alpha-Tubulin Mouse monoclonal antibody (dilution 1:400) by incubation with Alexa488 conjugated Goat Anti-Mouse IgG (H+L) antibody (dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Confocal imaging of NIH/3T3 cells using 67kDa Laminin Receptor Rabbit monoclonal antibody (STJ11103600, dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (dilution 1:500) (Red). The cells were counterstained with Alpha-Tubulin Mouse monoclonal antibody (dilution 1:400) by incubation with Alexa488 conjugated Goat Anti-Mouse IgG (H+L) antibody (dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Western blot analysis of various lysates using 67kDa Laminin Receptor Rabbit monoclonal antibody (STJ11103600) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.