

## Anti-TLR6 antibody (697-796) (STJ11103621)

## **GENERAL INFORMATION**

Product Type Primary antibodies

**Short Description** 

Applications WB/IF/ICC/ELISA Host/Source Rabbit Reactivity Human/Mouse

## **PRODUCT PROPERTIES**

Clonality Polyclonal Clone ID

Concentration Lot specific Conjugation Unconjugated Purification Affinity purification Dilution Range WB:1:500-1:1000

IF/ICC:1:50-1:200

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.

Isotype IaG

Storage Instruction Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

## **TARGET INFORMATION**

Gene ID 10333

Gene Symbol TLR6


Uniprot ID TLR6\_HUMAN

Immunogen


Immunogen 697-796

Region
Specificity A synthetic peptide corresponding to a sequence within amino acids 697-796 of human TLR6 (NP\_006059.2). KSIFVLSPNFVQSEWCHYEL YFAHHNLFHEGSNNLILILL EPIPQNSIPNKYHKLKALMT QRTYLQWPKEKSKRGLFWAN Immunogen

Sequence IRAAFNMKLTLVTENNDVKS



Western blot analysis of various lysates using TLR6 Rabbit pAb (STJ11103621) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (STJS000856) at 1:1000 dilution. Lysates/protiens: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit



Immunofluorescence analysis of Jurkat cells using TLR6 Rabbit pAb (STJ11103621) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit [36 (H+L) (STJ3001166) at 1:500 dilution. Blue: DAPI for nuclear stainino.