Anti-MANF antibody (25-182) [S4168RM] (STJ11104168)

STJ11104168

GENERAL INFORMATION

Product Type	Primary antibodies
Short Description	
Applications	WB/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES

Clonality	Monoclonal
Clone ID	S4168RM
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.05% Proclin300, 0.05% BSA, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	7873
Gene Symbol	MANF
Uniprot ID	MANF_HUMAN
Immunogen	
Immunogen Region	25-182
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 25-182 of human MANF (NP_006001.5).
Immunogen Sequence	LRPGDCEVCISYLGRFYQDL KDRDVTFSPATIENELIKFC REARGKENRLCYYIGATDDA ATKIINEVSKPLAHHIPVEK ICEKLKKKDSQICELKYDKQ IDLSTVDLKKLRVKELKKIL DDWGETCKGCAEKSDYIRKI NELMPKYAPKAASARTDL



Western blot analysis of extracts of various cell lines, using ARMET/ARP/MANF antibody (STJ11104168) at1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.



Western blot analysis of extracts of various cell lines, using ARMET/ARP/MANF antibody (STJ11104168) at1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.

St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081