

## Anti-CUL1 antibody (435-593) (STJ11104479)

STJ11104479

### GENERAL INFORMATION

<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	
<b>Applications</b>	WB/IHC-P/ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human/Mouse/Rat

### PRODUCT PROPERTIES


<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	Lot specific
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purification
<b>Dilution</b>	WB:1:500-1:1000
<b>Range</b>	IHC-P:1:50-1:200
	ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
<b>Formulation</b>	PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
<b>Isotype</b>	IgG
<b>Storage</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
<b>Instruction</b>	

### TARGET INFORMATION


<b>Gene ID</b>	8454
<b>Gene Symbol</b>	CUL1
<b>Uniprot ID</b>	CUL1_HUMAN
<b>Immunogen</b>	
<b>Immunogen Region</b>	435-593
<b>Specificity</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 435-593 of human Cullin 1 (NP_003583.2).
<b>Immunogen Sequence</b>	KNPEEALEDLTNQVMVFK YIEDKDVQKFYAKMLAKRL VHQNSASDDAEASMISKLKQ ACGFEYTSKLQRMFQDIGVS KDLNEQFKKHLTNSEPLDLD FSIQVLSSGSWPFQQSCTFA LPSELSYQRFTAFYASRH SGRKLTWLYQLSKGELVTN




Immunohistochemistry analysis of paraffin-embedded rat lung using Culin 1 Rabbit polyclonal antibody (STJ11104479) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded human lung cancer using Culin 1 Rabbit polyclonal antibody (STJ11104479) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded human colon using Culin 1 Rabbit polyclonal antibody (STJ11104479) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of various lysates, using CUL1 antibody (STJ11104479) at 1:500 dilution. Secondary antibody: HRP Coated Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081