

## Anti-SIRT5 antibody (216-310) (STJ11104572)

STJ11104572

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	
<b>Applications</b>	WB/ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human

### PRODUCT PROPERTIES

<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	Lot specific
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purification
<b>Dilution Range</b>	WB:1:100-1:500 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
<b>Formulation</b>	PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
<b>Isotype</b>	IgG
<b>Storage Instruction</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

### TARGET INFORMATION

<b>Gene ID</b>	23408
<b>Gene Symbol</b>	SIRT5
<b>Uniprot ID</b>	SIR5_HUMAN
<b>Immunogen</b>	
<b>Immunogen Region</b>	216-310
<b>Specificity</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 216-310 of human SIRT5 (NP_036373.1).
<b>Immunogen Sequence</b>	LRPHVWVFGENLDPAILIEEV DRELAHCDLCLVGTSSVVY PAAMFAPQVAARGVPVAEFN TETTPATNRFRHFQGPCGT TLPEALACHENETVS



Western blot analysis of extracts of various cell lines, using SIRT5 antibody (STJ11104572) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 180s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081