

Anti-CD34 antibody (160-320) (STJ11104581)

STJ11104581

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB:1:500-1:1000
Range	IF/CC:1:50-1:200
	ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	947
Gene Symbol	CD34
Uniprot ID	CD34_HUMAN
Immunogen	
Immunogen Region	160-320
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 160-320 of human CD34 (NP_001020280).
Immunogen Sequence	PYTSSSPILSDIAEKICSG IREVKLTQGICLEQNKTSSC AEFKKDRGEGLARVLCGEEQ ADADAGAQCSSLQAQSEVR PQCLLLVLNARTEISSKLQL MKKHQSDLKKGILDFTEQD VASHQSYSQKTLIALVTSGA LLAVLGITGYFLMNRSSWSP T



Immunofluorescence analysis of TF-1 cells using CD34 Rabbit polyclonal antibody (STJ11104581) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using CD34 Rabbit polyclonal antibody (STJ11104581) at 1:1000 dilution. Secondary antibody:HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection:ECL Basic Kit. Exposure time: 90s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081