

Anti-GSTA4 antibody (1-100) [S5149RM] (STJ11105149)

ST.I11105149

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications WB/ELISA Host/Source Rabbit Reactivity Human

PRODUCT PROPERTIES

Clonality Monoclonal S5149RM

Concentration Lot specific Unconjugated Purification

Purification

Dilution Range WB:1:500-1:1000

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.

Isotype IgG

Storage Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

Instruction

TARGET INFORMATION

Gene ID 2941 Gene Symbol GSTA4


Uniprot ID GSTA4_HUMAN

Immunogen Immunogen 1-100 Region

Specificity Recombinant funsion protein containing a sequence corresponding to amino acids 1-100 of human GSTA4 (NP_001503.1).

Immunogen MAARPKLHYPNGRGRMESVR WVLAAAGVEFDEEFLETKEQ LYKLQDGNHLLFQQVPMVEI DGMKLVQTRSILHYIADKHN

Sequence LFGKNLKERTLIDMYVEGTL



western blot analysis of lysates from HepCi2 Cells using GSTA4 Rabbit monoclonal antibody (STJ11105149) at 1:800 dilution. Secondary antibody HRP Goat Anti-Rabbit 1gG (H±L) (STJS000856) a 1:10000 dilution. Lysates/proteins: 25 Mu. g per lane Blocking buffer: 3% norlat dry milk in TBST. Detection ECL Basic Kft. Exposure time: 90s. western blot arraysis of rysales iron 2537 ceiss, using GSTA4 Rabbit monoclonal antibody (STJ11105149) at 1:800 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H-L) (STJS000556) at 1:1000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer; 3% nonfat dry milk: nT BST. Detection: ECL Basic Kit. Exposure time: 30s.