

Anti-BAD antibody (68-168 aa) (STJ11105422)

STJ11105422

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications WB/ELISA
Host/Source Rabbit
Reactivity Human/Mouse

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID

Concentration Lot specific
Conjugation Unconjugated
Purification Affinity purification
Dilution Range WB:1:500-1:1000

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.

Isotype IgG

Storage Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

Instruction

TARGET INFORMATION

Gene ID 572

Gene Symbol BAD Uniprot ID BAD_HUMAN

Immunogen


Immunogen 68-168 aa

Region


Specificity A synthetic peptide corresponding to a sequence within amino acids 68-168 of human Bad. (NP_004313.1).

Immunogen EIRSRHSSYPAGTEDDEGMG EEPSPFRGRSRSAPPNLWAA QRYGRELRRMSDEFVDSFKK GLPRPKSAGTATQMRQSSSW

Sequence TRVFQSWWDRNLGRGSSAPS Q



Western blot analysis of lysates from wild type (WT) and Bad knockdown (KD) 293T (KD) cells, using Bad Rabbit polyclonal antibody (STJ11105422) at 1:500 dilution. Secondary antibody: HRP Goat Arti-Rabbit IgG (H+L (STJS00856) at 1:1000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk ir TBST. Detactions: ECI Repic (Ht Exposure times 1806.



Western blot analysis of various lysates, using Bac Rabbit polyclonal antibody (STJ11105422) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbi IgG (H+L) (STJS000856) at 1:10000 dilution Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry, milk in TBST. Detection: ECL Basic Kt.