

## Anti-Phospho-LIPE-S660 antibody (STJ11105787)

STJ11105787

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	
<b>Applications</b>	WB/ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human/Mouse

### PRODUCT PROPERTIES

<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	Lot specific
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purification
<b>Dilution Range</b>	WB:1:500-1:1000 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
<b>Formulation</b>	PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
<b>Isotype</b>	IgG
<b>Storage Instruction</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

### TARGET INFORMATION

<b>Gene ID</b>	3991
<b>Gene Symbol</b>	LIPE
<b>Uniprot ID</b>	LIPS_HUMAN
<b>Immunogen</b>	QTSRS
<b>Immunogen Region</b>	
<b>Specificity</b>	A synthetic phosphorylated peptide around S660 of human LIPE (NP_005348.2).
<b>Immunogen Sequence</b>	QTSRS



Western blot analysis of lysates from 3T3-L1 cells, using Phospho-HSL-S660 Rabbit pAb (STJ11105787) at 1:600 dilution. 3T3-L1 cells were treated with Isoproterenol (10  $\mu$ M for 20 min) at 37  $^{\circ}$ C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25  $\mu$ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081