

Anti-Phospho-EIF4EBP1-T70 antibody [S5824RM] (STJ11105824)

ST.I11105824

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications WB/ELISA
Host/Source Rabbit
Reactivity Human/Mouse

PRODUCT PROPERTIES

Clonality Monoclonal
Clone ID 55824RM

Concentration Lot specific
Conjugation Purification
Dilution Range WB:1:1000-1:5000

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.05% Proclin300, 0.05% BSA, 50% Glycerol, pH 7.3.

Isotype IgG

Storage Instruction Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID 1978
Gene Symbol EIF4EBP1
Uniprot ID 4EBP1_HUMAN
Immunogen TKTPPR

Immunogen Region


Specificity A synthetic phosphorylated peptide around T70 of human eIF4EBP1 (NP_004086.1).

Immunogen TKTPPF

Sequence



Western blot analysis of extracts of NIH/3T3, usin Phospho-elf-4EBP1-TT0 antibody (STJ1110582 att:5000 dilution. NIH/3T3 cells were treated by Cl (20uL/400ul) at 37 ŰC for 1 hour. Secondary antibod HRP Goat Anti-Habbit IgG (H-L) (STJ3003656) 1:10000 dilution. Lysates/proteins: 25 Mu g per lan Blocking buffer 3% norlat dry milk in TBST. Detection



western bloch analysis of Wardsch or Archius Gellanse at 1500 dilution. 293T cells were treated by 10% FB at 1500 dilution. 293T cells were treated by 10% FB at 37 A°C for 30 minutes after serum-starvatio overnight. NH/13T3 cells were treated by Cli (20uL/40ul) at 37 A°C for 1 hour. Secondary antibod. HPP Goat Ann-Rabbit IgG (H-I), STL5000856) at 11:10000 dilution. Lystensephreties: 25 Mig per land. 11:10000 dilution. Lystensephreties: 25 Mig per land. 15:10000 dilution. Lystensephreties: 27 Mig per land.