

Anti-SP1 antibody (581-680 aa) (STJ11107080)
STJ11107080

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	IHC-P/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage	
Instruction	

TARGET INFORMATION


Gene ID	6667
Gene Symbol	SP1
Uniprot ID	SP1_HUMAN
Immunogen	
Immunogen Region	581-680 aa
Specificity	A synthetic peptide corresponding to a sequence within amino acids 581-680 of human SP1 (NP_612482.2).
Immunogen Sequence	GGEEGENSPDAQPQAGRRTREACTCPYCKDSEGRGSGDP GKKKQHIHQGCGKVYGT SHLRAHLRWHTGERPFMCTW SYCGKRFRTRSDQLQRHKRTH




Immunohistochemistry analysis of SP1 in paraffin-embedded human small intestine tissue using SP1 Rabbit pAbA (STJ11107080) at a dilution of 1:200 (40x lens). A High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to immunohistochemistry staining.



Immunohistochemistry analysis of SP1 in paraffin-embedded human breast tissue using SP1 Rabbit pAbA (STJ11107080) at a dilution of 1:200 (40x lens). A High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to immunohistochemistry staining.



Immunohistochemistry analysis of SP1 in paraffin-embedded human spleen tissue using SP1 Rabbit pAbA (STJ11107080) at a dilution of 1:200 (40x lens). A High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to immunohistochemistry staining.



Immunohistochemistry analysis of SP1 in paraffin-embedded rat colon tissue using SP1 Rabbit pAbA (STJ11107080) at a dilution of 1:200 (40x lens). A High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to immunohistochemistry staining.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081