

Anti-CHRNA7 antibody (23-230) (STJ111098)

STJ111098

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB:1:500-1:2000
Range	IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
Instruction	

TARGET INFORMATION


Gene ID	1139/89832
Gene Symbol	CHRNA7
Uniprot ID	ACHA7_HUMAN
Immunogen	
Region	23-230
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 23-230 of human CHRNA7 (NP_001177384.1).
Immunogen Sequence	ASPPSTPPWDPGHIPGASVR PAPGPVSLQGEFQRKLYKEL VKNYNPLERPVANDSQPLTV YFSLSLQIMDVDEKNQVLT TNIWLQMSWTDHYLQWNVSE YPGVKTVRFDPGQIWKPDIL LYSADERFDATFHTNVLVN SSGHCQYLPPIGIFKSSCYID VRWFPPDVQHCKLKFGWSY GWSLDLQMQEADISGYIPN GEWDLVGI



Immunofluorescence analysis of A L-929 cells using Anti-CHRNA7 Rabbit polyclonal antibody (STJ111098) at a dilution of 1:200 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of CHRNA7 in paraffin-embedded human placenta using Anti-CHRNA7 Rabbit polyclonal antibody (STJ111098) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of lysates from MCF7 cells using Anti-CHRNA7 Rabbit polyclonal antibody (STJ111098) at a dilution of 1:2000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ5000856) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 120s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081