

Anti-HOXA10 antibody (236-336) (STJ111287)

STJ111287

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications WB/ELISA Host/Source Rabbit

Reactivity Human/Mouse/Rat

PRODUCT PROPERTIES

Clonality Polyclonal Clone ID

Dilution Range WB:1:500-1:2000

Concentration Lot specific
Conjugation Unconjugated
Purification Affinity purification

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.

Isotype IgG

Storage Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

Instruction

TARGET INFORMATION

Gene ID 3206
Gene Symbol HOXA10
Uniprot ID HXA10_HUMAN


Immunogen

Immunogen 236-336


Region

Specificity Recombinant fusion protein containing a sequence corresponding to amino acids 236-336 of human HOXA10 (NP_061824.3). **Immunogen** LGAGPFPAQPPGRGFDLPPA LASGSADAARKERALDSPPP PTLACGSGGGSQGDEEAHAS SSAAEELSPAPSESSKASPE


Sequence KDSLGNSKGENAANWLTAKS G




Western blot analysis of extracts of various cell lines using HOXA10 antibody (STJ111287) at 1:1000 dilution Secondary antibody: HRP Goat Anti-rabbit IgG (H-L) a 1:10000 dilution. Lysates/proteins: 25ug per label Blocking buffer: 3% non-fat dry milk in TBS1. Detection ECI. Resic Kit Expresser unes: 10s.



Western blot analysis of extracts of Mouse kidne using HOXA10 antibody (STJ111287) at 1:1000 dilution Secondary antibody: HRP Goat Anti-Rabbit IgG (H+ (STJS000856) at 1:10000 dilution. Lysates/proteins: 2 Mu g per lane. Blocking buffer: 3% nonfat dry milk i TBST. Detection: ECL Enhanced Kit. Exposure time



Western blot analysis of extracts of mouse kidne using HOXA/10 antibody (STJ111287) at 1:1000 dilutio Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) 1:10000 dilution. Lysates/proteins: 25ug per la Blocking buffer: 3% non-fat dry milk in TBST. Detectic ECI. Enhanced kif. Evonsure time: 100



Western blot analysis of extracts of various cell lines using HOXA10 antibody (STJ111287) at 1:1000 dillution Secondary antibody: HFP Goat Anti-Rabbit IgG (H-L (STJS000856) at 1:1000 dilution. Lysates/proteins (STJS000856) at 1:1000 dilution. Lysates/proteins Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST_Dates/critica FC/I Barsic ME Exerceptus/time; Vis-