

## Anti-VAMP1 antibody (1-96) (STJ111459)

STJ111459

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	
<b>Applications</b>	WB/IHC-P/ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human/Mouse/Rat

### PRODUCT PROPERTIES


<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	Lot specific
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purification
<b>Dilution Range</b>	WB:1:500-1:2000 IHC-P:1:50-1:100 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
<b>Formulation</b>	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
<b>Isotype</b>	IgG
<b>Storage Instruction</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

### TARGET INFORMATION


<b>Gene ID</b>	6843
<b>Gene Symbol</b>	VAMP1
<b>Uniprot ID</b>	VAMP1_HUMAN
<b>Immunogen</b>	
<b>Immunogen Region</b>	1-96
<b>Specificity</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 1-96 of human VAMP1 (NP_954740.1).
<b>Immunogen Sequence</b>	MSAPAQPPAEGTEGTAPGGG PPGPPNMTSNRRLQQTQQA VEEVDIIRVNVDKVLERDQ KLSLDDRADALQAGASQFE SSAAKLKRKYWWKNCK




Western blot analysis of extracts of various cell lines, using VAMP1 antibody (STJ111459) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 5s.



Immunohistochemistry analysis of paraffin-embedded mouse lung using VAMP1 antibody (STJ111459) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry of paraffin-embedded rat ovary using VAMP1 antibody (STJ111459) at dilution of 1:100 (40x lens).



Immunohistochemistry analysis of paraffin-embedded human gastric cancer using VAMP1 antibody (STJ111459) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.