

Anti-MARCO antibody (421-520) (STJ112088)

STJ112088

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
Instruction	

TARGET INFORMATION


Gene ID	8685
Gene Symbol	MARCO
Uniprot ID	MARCO_HUMAN
Immunogen	
Immunogen Region	421-520
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 421-520 of human MARCO (NP_006761.1). SVSVRIVGSSNRGAEVYYS GTWGTICDDEWQNSDAIVFC RMLGYSKGRALYKVGAGTGQ IWLDNVQCRGTESTLWSC TK
Immunogen Sequence	NSWGHHDCSHEEDAGVECSV



Immunofluorescence analysis of mouse lung cells using MARCO Rabbit polyclonal antibody (STJ112088) at dilution of 1:200 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of rat lung cells using MARCO Rabbit polyclonal antibody (STJ112088) at dilution of 1:200 (40x lens). Blue: DAPI for nuclear staining.



Western blot analysis of extracts of various cell lines, using MARCO antibody (STJ112088) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.