

Anti-KIF14 antibody (1399-1648) (STJ112313)

STJ112313

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB:1:1000-1:2000
Range	IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
Instruction	

TARGET INFORMATION


Gene ID	9928
Gene Symbol	KIF14
Uniprot ID	KIF14_HUMAN
Immunogen	
Region	1399-1648
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1399-1648 of human KIF14 (NP_055690.1).
Immunogen Sequence	LENGNNKAASVQEEFMDAVC DGVGLGMKILLDSGLEKAKE LQHELFRQCTKNEVTKEMKT NAMGLIRSLNIFAESKIKS FRRQVQEEFQDFKRMVN RAPEFLKLKHCLEKAEIIL SALKGCHSDINLLQTCVESI RNLASDFYSDFSVPSTSVGS YESRVTHIVHQLLESLSL LFCFESEESPDLLKPWETYN QNTKEEHQQSKSSGIDGSKN KGVPKRVYELHGSSPAVSS




Immunofluorescence analysis of U-2 OS cells using KIF14 Polyclonal Antibody (STJ112313) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of L929 cells using KIF14 Polyclonal Antibody (STJ112313) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded human breast cancer using KIF14 antibody (STJ112313) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat ovary using KIF14 antibody (STJ112313) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081