

Anti-DAB1 antibody (1-555) (STJ112387)

STJ112387

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB:1:100-1:500
Range	IF/CC:1:50-1:200
	ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	1600
Gene Symbol	DAB1
Uniprot ID	DAB1_HUMAN
Immunogen	
Immunogen Region	1-555
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-185 of human DAB1 (NP_066566.3).
Immunogen Sequence	MSTETELQVAVKTSAKKDSR KKGQDRSEATLKRFKGEV RYKAKLIGIDEVSAARGDKL CQDSMMKLKGVVAGARSKGE HKQKIFLTISFGGIKIFDEK TGAHQHHAVHEISYIAKDI TDHRAFGYVCGKEGNHRFVA IKTAQAAEPVILDLRDLFQL IYELKQREELEKKAQDKQC EQAVY



Immunofluorescence analysis of SH-SY5Y cells using DAB1 Rabbit polyclonal antibody (STJ112387) at dilution of 1:200 (40x lens). Blue: DAPI for nuclear staining.



Western blot analysis of extracts of various cell lines, using DAB1 antibody (STJ112387) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ3000896) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081